



The Weakly Bound 1:1 Complex of CO₂ and H₂O

Observation and Assignment of Intermolecular van der Waals Vibrations

Andersen, Jonas; Heimdal, J.; Wallin Mahler Andersen, Denise; Nelander, B.; Larsen, René Wugt

Published in:
MAX-lab Activity Report 2012

Publication date:
2013

[Link back to DTU Orbit](#)

Citation (APA):
Andersen, J., Heimdal, J., Wallin Mahler Andersen, D., Nelander, B., & Larsen, R. W. (2013). The Weakly Bound 1:1 Complex of CO₂ and H₂O: Observation and Assignment of Intermolecular van der Waals Vibrations. In *MAX-lab Activity Report 2012*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The Weakly Bound 1:1 Complex of CO₂ and H₂O: Observation and Assignment of Intermolecular van der Waals Vibrations

J. Andersen^a, J. Heimdal^b, B. Nelander^b, D.W. Mahler^a and R. Wugt Larsen^a

^a*Technical University of Denmark, Department of Chemistry, DK-2800 Kongens Lyngby, Denmark*

^b*Lund University, MAX-IV Laboratory, SE-22100 Lund, Sweden*

The intermolecular interaction between CO₂ and H₂O plays a major role for a multifaceted variety of phenomena in physics, chemistry and biology including the radiation transfer through the Earth's and planetary atmospheres, the early stage of carbonic acid formation and the transport of dissolved CO₂ in the tissues of biological organisms.

The prototypical binary mixed 1:1 complex formed between CO₂ and H₂O has accordingly been investigated intensively by both theory and experiment. A recent comprehensive work reporting a complete high-level 5-D *ab initio* intermolecular potential energy surface composed of 23 000 single-point energies indicates that the global potential energy minimum has a planar and T-shaped geometry of C_{2v} symmetry with the oxygen atom of H₂O bound to the C atom and the H atoms pointing away from the CO₂ molecule as shown in Fig. 1.¹

The weakly bound nature of the CO₂–H₂O system results in rather small perturbations of the intramolecular vibrational transitions relative to the vibrational transitions of the isolated subunits. The total shift of the zero-point energy contribution to D_0 caused by perturbations of intramolecular vibrational transitions is in the order of 5 cm⁻¹. A reliable determination of the system's zero-point energy thus relies completely on accurate band origins for the set of five intermolecular vibrational transitions introduced by the van der Waals interactions.¹

Terahertz absorption spectra have recently been obtained at the infrared beam-line for CO₂/H₂O mixtures embedded in cryogenic neon matrices at 2.8 K. Several millimeter thick matrices of neon doped with CO₂ and H₂O were deposited at 3.6 K on a gold-plated OFHC copper mirror inside an immersion helium cryostat (IHC-3) modified for matrix isolation spectroscopy.² Single-beam spectra were collected by the Bruker IFS 120 FTIR spectrometer located at the beam-line.

Based upon relative concentration dependency measurements, heavy isotope substitutions and complementary mid-infrared studies of perturbed intramolecular vibrational transitions, two high-frequency intermolecular vibrational transitions have been observed for the H₂O–CO₂ van der Waals complex at 101.8 and 166.2 cm⁻¹ as shown in Figure 2. These large-amplitude

intermolecular modes have been assigned to out-of-plane wagging and in-plane rocking of the H₂O subunit, respectively, and provide crucial observables for benchmark theoretical descriptions of this van der Waals system's flat intermolecular potential energy surface.




Figure 1: The five intermolecular coordinates (r , θ' , α' , β' , γ') specifying the configuration of the weakly bound H₂O–CO₂ van der Waals complex. The white circles represent oxygen atoms and the grey circles carbon and hydrogen.




Figure 2: The assigned transitions for the recorded THz spectra of CO₂/H₂O mixtures embedded in solid neon matrices at 2.8 K.

REFERENCES

- [1] J. Makarewicz, *Journal of Chemical Physics*, 2010, **132**, 234305
- [2] J. Ceponkus, P. Uvdal, and B. Nelander, *Journal of Chemical Physics*, 2013, **138**, 244305